When used as directed, this product is effective against a wide variety of ‡ pathogenic microorganisms. At a 1:64 dilution (2 oz. of product per gallon of water), in the presence of 200 ppm hard water, 5% serum load and 10 minute contact time, unless otherwise noted, this product kills the following on hard, non-porous inanimate surfaces:

‡Bactericidal Activity: Pseudomonas aeruginosa, Staphylococcus aureus, Salmonella enterica, Escherichia coli (0157:H7), Streptococcus pneumoniae.

Antibiotic-Resistant Bactericidal Activity: Staphylococcus aureus. Methicillin resistant (MRSA), Staphylococcus aureus. Community Associated Methicillin-resistant (CA-MRSA). (Genotype USA300). Staphylococcus aureus. Community Associated Methicillin-resistant (CA-MRSA), (Genotype USA400), Staphylococcus epidermidis. Methicillin-resistant (MRSE). Streptococcus pneumoniae, Penicillin-resistant (PRSP), ***Enterococcus faecium (Vancomycin Resistant (VRE)).

***Indicates that a dilution of 4 oz. per gallon of water is required for this claim.

**Virucidal Activity: *Adenovirus type 8 (5 minute contact time), *Hepatitis B Virus (HBV) (Duck Hepatitis B Virus as the surrogate) (5 minute contact time on pre-cleaned surfaces), *Hepatitis C Virus (HCV) (Bovine Viral Diarrhea Virus as the surrogate) (5 minute contact time), *Herpes Simplex Virus Type 1 (5 minute contact time), *Herpes Simplex Virus Type 2 (5 minute contact time), Kills *HIV-1 (Human Immunodeficiency Virus Type 1) (AIDS virus) (1 minute contact time), *Influenza Virus Type A (H1N1) (1 minute contact time), *Norovirus (Feline Calicivirus as the surrogate) (5 minute contact time), *Rhinovirus Type 37 (5 minute contact time), *Rotavirus (5 minute contact time), +*SARS-Related Coronavirus 2 (SARS CoV-2), BEI Resources NR-52281, Strain Isolate USA-WA1/2020 (the virus that causes COVID-19) (1 minute contact time).

††Indicates testing in the presence of 400 ppm hard water.

Modes of Application: This product can be applied by mop, sponge, cloth, low pressure mechanical spray device, coarse trigger sprayer, or auto-scrubber. Change cloth, sponges or towels frequently to avoid redeposition of soil.

Fungicidal Performance: This product kills the following fungi in 10 minutes at 1:16 (8 oz. per gallon of water). in the presence of 200 ppm hard water. 5% serum load, on hard. non-porous surfaces: Trichophyton interdigitale (Athlete's foot fungus).

DIRECTIONS FOR USE: It is a violation of Federal Law to use this product in a manner inconsistent with its labeling.

This product is not to be used as a terminal sterilant/high level disinfectant on any surface or instrument that (1) is introduced directly into the human body, either into or in contact with the blood stream or normally sterile areas of the body, or (2) contacts intact mucous membranes but which does not ordinarily penetrate the blood barrier or otherwise enter normally sterile areas of the body. This product may be used to pre-clean or decontaminate critical or semi-critical medical devices prior to sterilization or high level disinfection. Not recommended for use on copper, brass, granite, marble or zinc.

To Prepare Use Solution:

Add the product at 2 oz. of product per gallon of water (1:64).

For Use as a One-Step Cleaner/Disinfectant: Dilute at 2 oz. of product per gallon of water (1:64), Pre-clean visibly soiled surfaces. Apply Use Solution with a sponge, brush, cloth. mop, low pressure mechanical spray device, or coarse trigger sprayer to hard, non-porous surfaces. Spray 6 - 8 inches from surface, making sure to wet surfaces thoroughly. All surfaces must remain visibly wet for 10 minutes. Wipe surfaces or let air dry.

Use 4 oz. per gallon of water to kill Enterococcus faecium (Vancomycin Resistant (VRE).

NOTE: Rinsing is not necessary unless floors are to be coated with finish or restorer. When cleaning floors position "wet floor" signs around area to be cleaned. Floors will be slippery when wet or contaminated with foreign materials. Do not use on glassware, utensils, or dishes.

For use as a *Virucide: Dilute at 2 oz. of product per gallon of water (1:64). Pre-clean visibly soiled surfaces. Apply Use Solution until thoroughly wet to hard, non-porous surfaces. All surfaces must remain visibly wet for 5 minutes. A one minute contact time is required for *HIV-1 (AIDS virus). *Influenza Virus Type A (H1N1). *++SARS-Related Coronavirus 2 (the virus that causes COVID-19). Wipe surfaces or let air dry.

This product kills HIV-1 (associated with AIDS), HBV and HCV on pre-cleaned surfaces/object previously soiled with blood and body fluids.

On pre-cleaned surfaces/objects previously soiled with blood/body fluids in health care settings or other settings in which there is an expected likelihood of soiling of inanimate surfaces/ objects with blood or body fluids and in which the surfaces/objects likely to be soiled with blood or body fluids can be associated with the potential for transmission of Human Immunodeficiency Virus Type 1 (HIV-1) (associated with AIDS), Hepatitis B Virus (HBV), or Hepatitis C Virus (HCV),

SPECIAL INSTRUCTIONS FOR CLEANING AND DECONTAMINATION AGAINST HBV, HCV and HIV-1 OF SURFACES/OBJECTS SOILED WITH BLOOD/BODY FLUIDS. Personal Protection: Disposable latex or vinyl gloves, gowns, face masks, and eye coverings must be worn during all cleaning of body fluids, blood, and decontamination procedures. Cleaning Procedures: Blood and body fluids must be thoroughly cleaned from surfaces and objects before application of this product. Contact Time: Allow surface to remain wet for 1 minute to kill HIV-1.5 minutes to kill HBV & HCV. Disposal of Infectious Material: Blood and other body fluids must be autoclayed and disposed of according to Federal. State, and local regulations for infectious waste disposal.

To Kill Fungi: Pre-clean visibly soiled areas, Prepare use solution by adding 8 oz. of this product per gallon of water (1:16). Apply use solution to hard, non-porous surfaces. Allow surface to remain visably wet for 10 minutes. Wipe surfaces or let air dry.

Animal Premise Virucidat* Performance: This product kills the following viruses in 10 minutes at 1:16 (8 oz. per qal. of water) and 5% soil on hard, non-porous surfaces: *Canine Parvovirus (CPV)

Peroxide **Disinfectant Cleaner**

Victoria Bay

For Use as a One-Step Cleaner/Disinfectant:

Wear chemical splash-proof goggles or face





Dilute at 2 oz. of product per gallon of water



Spray surface until thoroughly wet.





Allow surface to remain wet. Read label for contact times.



Wipe surface and let air dry.

Read label for entire directions for use.

Peroxide **Disinfectant Cleaner** Victoria Bay

Effective against SARS-CoV-2 in 1 minute on hard, nonporous surfaces.

ACTIVE INGREDIENT:

REORDER #

D02030

Hydrogen Peroxide	4.25%
OTHER INGREDIENTS:	<u>95.75%</u>
TOTAL:	100.00%

KEEP OUT OF REACH OF CHILDREN CAUTION

See additional precautionary statements on side.

FIRST AID

IF IN EYES: Hold eve open and rinse slowly and gently with water for 15-20 minutes. Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye. IF ON SKIN OR CLOTHING: Take off contaminated clothing. Rinse skin immediately with plenty of water for 15-20 minutes. **IF SWALLOWED:** Have person sip a glass of water if able to swallow. Do not induce vomiting unless told to do so by a poison control center or doctor. Do not give anything by mouth to an unconscious

IN CASE OF EMERGENCY, CALL A POISON CONTROL CENTER OR DOCTOR FOR TREATMENT ADVICE. Have the product container or label with you when calling a Poison Control Center or doctor or going in for treatment. Note to Physician: Probable mucosal damage may contraindicate gastric

> Manufactured for: **Victoria Bay Products** 255 Route 1 & 9 Jersey City, NJ 07306 In case of emergency call (800) 535-5053



For Treatment of Animal Housing Facilities: Dilute at 2 oz. of product per gallon of water (1:64).† Remove all animals and feeds from areas being treated. Remove all litter and manure from floors, walls and surfaces of barns, pens, stalls, chutes, and other facilities occupied or traversed by animals. Empty or cover all troughs, racks and other feeding and watering appliances. Thoroughly clean all surfaces with soap or detergent and rinse with water. Apply fresh Use Solution to floors. walls, cages and other washable hard, non-porous surfaces. For smaller surfaces, use a trigger spray bottle to spray all surfaces with solution until wet. To disinfect, all surfaces must remain visably wet for ten minutes. Immerse all halters, ropes, and other types of equipment used in handling and restraining animals, as well as forks, shovels, and scrapers used for removing litter and manure. Ventilate buildings and other closed spaces. Do not house animals or re-employ equipment until product has dried. For disinfection of feed racks, troughs, automatic feeders. fountains and watering appliances scrub with use-solution, surfaces must remain visibly wet for 10 minutes. Then thoroughly scrub all treated surfaces with soap or detergent and rinse with potable water before reuse.

†Prepare a Use Solution of 8 oz. of product per gallon of water for effectiveness against Canine Parvovirus (CPV) (1:16).

Sanitizing: When used as a non-food contact sanitizer at a 1:128 dilution (1 oz. of product per gallon of water), in the presence of 200 ppm hard water, 5% serum load, this product reduces the following bacteria by 99.9% with a 5 minute contact time on hard non-porous inanimate surfaces: Enterobacter aeroaenes, Salmonella enterica

To Sanitize Non-Food Contact Surfaces: Dilute at 1 oz. of product per gallon of water (1:128). Apply Use Solution until thoroughly wet. All surfaces must remain visibly wet for 5 minutes Wipe surfaces or let air dry. Not for use on food contact surfaces or on food preparation areas. NOTE: Do not use on glassware, utensils, or dishes.

For Use as a Deodorizer: Dilute product at 0.5-1 oz. of product per gallon of water (1:256-1:128). Apply to hard, non-porous surfaces. Let stand for 10 minutes. Wipe surfaces or let air dry.

For Use as a Glass Cleaner: Dilute product at 1 oz. of product per gallon of water (1:128). Apply evenly to glass surfaces through a coarse trigger spray device. Do not breathe spray. Wipe dry with an absorbent cloth or towel.

PRECAUTIONARY STATEMENTS

HAZARDS TO HUMANS AND DOMESTIC ANIMALS

CAUTION: Harmful if swallowed. Causes moderate eye irritation. Avoid contact with eyes or clothing. Wear chemical splash-proof goggles or face shield. Wash thoroughly with soap and water after handling and before eating, drinking, chewing gum, using tobacco, or using the toilet.

STORAGE AND DISPOSAL

Do not contaminate water, food or feed by storage or disposal.

PESTICIDE STORAGE: Do not reuse empty container. PESTICIDE DISPOSAL: Pesticide wastes are acutely hazardous. Improper disposal of excess pesticide, spray mixture, or rinsate is a violation of Federal law. If these wastes cannot be disposed of by use according to label instructions, contact your State Pesticide or Environmental Control Agency, or the Hazardous Waste representative at the nearest EPA Regional Office for guidance. **CONTAINER DISPOSAL:** Non-refillable container. Do not reuse this container to hold materials other than pesticides or diluted pesticide rinsate. Triple rinse as follows: Fill container ¼ full with water and recap. Shake for 10 seconds. Drain for 10 seconds after the flow begins to drip. Follow Pesticide Disposal instructions for rinsate disposal. Repeat procedure two more times. Then offer for recycling or reconditioning if appropriate or puncture and dispose of in a sanitary landfill or by incineration.

EPA REG. NO. 45745-11-68168, EPA EST. NO. 45745-TN-2

